was used in paint, gasoline, water pipes, and many other products. Now that we know the dangers of lead, house paint is almost lead-free, leaded gasoline has been phased out, and household plumbing is no longer made with lead materials.

Children

NOY buA

Environmental Protection

Agency

United States

ead Poisoning

October 2000

EPA 747-K-00-003

children

Office of Pollution Prevention and Toxics (7404)

Lead awareness and your

About 1 in 22 children in America have

high levels of lead in their blood, according to

the Centers for Disease Control and Prevention. You may have lead around your building without

knowing it because you can't see, taste, or smell

lead. You may have lead in the dust, paint, or soil

in and around your home, or in your drinking water

or food. Because it does not break down naturally,

lead can remain a problem until it is removed.


Before we knew how harmful it could be, lead

How lead affects your

Children who should be tested under your state or local screening plan.

To find out where to have your child tested, call your doctor or local health clinic. They can explain what the test results mean, and if more testing will be needed.

2. Keep it clean.



- peeling or chipping paint are dangerous if eaten. Lead paint in good condition is not usually a problem except in places where painted surfaces rub against each other and create dust. (For example, when you open a window, the painted surfaces rub against each other.)
- 3. Reduce the risk from lead paint.
- Make sure your children wash their hands before meals, nap time, and bedtime.

Most homes built before 1960 contain

leaded paint. Some homes built as recently as

1978 may also contain lead paint. This paint

could be on window frames, walls, the outside

of your house, or other surfaces. Tiny pieces of

lead paint yourself. thoroughly. Don't try to remove the proper equipment to clean up

oud a thorough cleanned is done. the building until all work is finished and pregnant women, should leave All occupants, especially children

into your home. 5. Don't bring lead dust

that used leaded gas. from years of exhaust tumes from cars and trucks Soil by roads or highways may be contaminated from lead paint on the outside of the building. Soil very close to homes may be contaminated tracking in lead from the soil around your home. home on your hands or clothes. You may also be lead, you may unknowingly bring lead into your shop or lead factory, or if your hobby involves painting, with batteries, or in a radiator repair It you work in construction, demolition or

- before you go home. hobby, change your clothes and shower If you work with lead in your job or
- come inside. sure they wash their hands when they your children from eating dirt, and make sticks to fingers and toys. Try to keep or grassy areas instead of dirt which Encourage your children to play in sand

drinking water. 6. Get lead out of your

through the skin. problem because lead does not enter the body reduce the amount of lead. Bathing is not a with lead materials. Boiling the water will not your home from household plumbing that is made contain lead. Water usually picks up lead inside Most well or city water does not naturally



3) Call EPA's Safe Drinking Water Hotline

drinking it, especially if you have not

1) Use only cold water for drinking, cooking,

it has sat for a long time in the pipes, is hot,

Household water will contain more lead it

supplier to see how to get it tested.

local health department or your water

your water is to have it tested. Call your

ui pool only way to know ij you have lead in

2) Run water for 15 to 30 seconds before

It you think your plumbing might have

used your water for a tew hours.

and making baby formula.

:11 ni bsəl

or is naturally acidic.

for more information.

7. Eat right.

are high in calcium. eggs, lean red meat, and beans. Dairy products will absorb less lead. Foods rich in iron include A child who gets enough iron and calcium

- glassware or imported or old pottery. Don't store food or liquid in lead crystal
- the bag. food, keep the printing on the outside of It you reuse plastic bags to store or carry

child's health

The long-term effects of lead in a child can be severe. They include learning disabilities, decreased growth, hyperactivity, impaired hearing, and even brain damage. If caught early, these effects can be limited by reducing exposure to lead or by medical treatment. If you are pregnant, avoid exposing yourself to lead. Lead can pass through your body to your baby. The good news is that there are simple things you can do to help protect your family.

1. Get your child tested.

Even children who appear healthy may have high levels of lead. You can't tell if a child has lead poisoning unless you have him or her tested. A blood test takes only ten minutes, and results should be ready within a week.

Blood tests are usually recommended for:

- Children at ages 1 and 2. 1
- Children or other family members who have been exposed to high levels of lead.

Ordinary dust and dirt may contain lead. Children can swallow lead or breathe lead contaminated dust if they play in dust or dirt and then put their fingers or toys in their mouths, or if they eat without washing their hands first.

- Keep the areas where your children play / as dust-free and clean as possible.
- / Wash pacifiers and bottles after they fall on the floor. Keep extras handy.
- Clean floors, window frames, window 1 sills, and other surfaces weekly. Use a mop, sponge, or paper towel with warm water and a general all-purpose cleaner or a cleaner made specifically for lead. REMEMBER: NEVER MIX AMMONIA AND BLEACH PRODUCTS TOGETHER SINCE THEY CAN FORM A DANGEROUS GAS.
- Thoroughly rinse sponges and mop heads after cleaning dirty and dusty areas.
- Wash toys and stuffed animals regularly.

Don't burn painted wood. It may contain 1 lead.

Make sure your child does not chew on

anything covered with lead paint, such as

painted window sills, cribs, or playpens.

4. Don't remove lead paint yourself

Families have been poisoned by scraping or sanding lead paint because these activities generate large amounts of lead dust. Lead dust from repairs or renovations of older buildings can remain in the building long after the work is completed. Heating paint may release lead into the air.

- Ask your local or state health department 1 if they will test your home for lead paint. Some will test for free. Home test kits cannot detect small amounts of lead under some conditions.
- 1 Hire a person with special training for correcting lead paint problems to remove lead paint from your home, someone who knows how to do this work safely and has

Protect Your Children From Lead Poisoning

Get your child tested for leadpoisoning, even if he or she seems healthy.


Clean floors, window frames, window sills, and other surfaces weekly. Use a mop, sponge, or paper towel with warm water and a general all-purpose cleaner or a cleaner made specifically for lead.


Reduce the risk of lead paint. Make sure your child is not chewing on anything covered with lead paint.

Don't try to remove lead paint yourself.

Don't bring lead dust into your home from work or a hobby.

> Have your water tested. If the cold water hasn't been used for more than a few hours, let it run for 15–30 seconds before drinking it or cooking with it.





Lead poisoning is a serious problem for young children the younger the child, the greater the risk.

For More Information

EPA's Safe Drinking Water Hotline 1-800-426-4791 National Lead Information Center 1-800-424-LEAD

Visit our web site http://www.epa.gov/lead

United States Environmental Protection Agency

Office of Pollution Prevention and Toxics

Office of Ground Water and Drinking Water

Recycled/Recyclable — Printed with vegetable oil based inks on recycled paper (30% minimum post-consumer)